

Homework 2

1. Let \hat{A}_1 , \hat{A}_2 and \hat{A}_3 be operators which are defined as

$$\hat{A}_1 f(t) = \frac{df(t)}{dt} + 3f(t) \quad (1)$$

$$\hat{A}_2 f(t) = \frac{d^2 f(t)}{dt^2} \quad (2)$$

$$\hat{A}_3 f(t) = \frac{d^2 f(t)}{dt^2} - 4k^2 t^2 f(t) \quad (3)$$

- (a) Show that all three operators are *linear*.
(b) Determine the eigenvalues of operators \hat{A}_1 and \hat{A}_2 and the corresponding eigenfunctions. Is the set of eigenvalues discrete or continuous in these two cases? Explain. (**Hint:** The easiest way to solve the first two problems is to use Laplace transforms. It is also helpful to consider the positive and negative eigenvalues of \hat{A}_2 separately).
(c) Show that

$$f(t) = e^{-kt^2} \quad (4)$$

is an eigenfunction of operator \hat{A}_3 , and determine the eigenvalue that corresponds to it (you can assume that k is given).

2. Show that functions $\{1, \cos x, \cos 2x, \dots, \cos nx, \dots\}$ are orthogonal with respect to scalar product

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f(x)g(x)dx \quad (5)$$

How should these functions be modified so that they become *orthonormal*?

3. Consider the sequence of functions $\{1, x, x^2, \dots, x^n, \dots\}$, which belong to a functional space where the scalar product is defined as

$$\langle f, g \rangle = \int_0^1 f(x)g(x)dx \quad (6)$$

- (a) Show that these functions are *not* orthogonal.
(b) Using set $\{1, x, x^2, \dots, x^n, \dots\}$ as the starting point, apply the Gram-Schmidt procedure to obtain a sequence of orthogonal polynomials $\{p_0(x), p_1(x), \dots, p_n(x), \dots\}$. For the purposes of this problem, it will suffice to explicitly construct only polynomials $p_0(x)$, $p_1(x)$ and $p_2(x)$. Show all your work!

(c) Are the polynomials obtained in this manner orthonormal, or just orthogonal? Explain.

4. Let S be a linear vector space in which functions ψ_0 and ψ_1 constitute an orthonormal basis, and suppose that operators \hat{A}_0 and \hat{A}_1 are defined as

$$\hat{A}_0\psi = \langle\psi_0, \psi\rangle\psi_0 \quad (7)$$

and

$$\hat{A}_1\psi = \langle\psi_1, \psi\rangle\psi_1 \quad (8)$$

- (a) Show that ψ_0 and ψ_1 are eigenfunctions of both \hat{A}_0 and \hat{A}_1 , and determine the corresponding eigenvalues.
- (b) Find the matrix representations of operators \hat{A}_0 and \hat{A}_1 in basis $\{\psi_0, \psi_1\}$.
5. Let S be a linear vector space in which functions ψ_0 and ψ_1 are an orthonormal basis, and suppose that operator \hat{A} transforms functions of the form

$$\psi = \alpha_0\psi_0 + \alpha_1\psi_1 \quad (9)$$

into

$$\hat{A}\psi = \frac{1}{\sqrt{2}}(\alpha_0 + \alpha_1)\psi_0 - \frac{1}{\sqrt{2}}(\alpha_0 - \alpha_1)\psi_1 \quad (10)$$

- (a) Find the eigenfunctions of this operator, and the eigenvalues that correspond to them. **Note:** Make sure that your eigenfunctions are *normalized*.
- (b) Show that the eigenfunctions you obtained in part (a) are orthogonal.
6. In this problem we will be interested in a pair of particles whose wave functions are

$$\psi_+ = \frac{1}{\sqrt{2}}\psi_0 + \frac{1}{\sqrt{2}}\psi_1 \quad (11)$$

and

$$\psi_- = \frac{1}{\sqrt{2}}\psi_0 - \frac{1}{\sqrt{2}}\psi_1 \quad (12)$$

respectively. When these two particles interact, we know that the resulting composite state will have the form

$$\Psi = \psi_+ \otimes \psi_- \quad (13)$$

where $\psi_+ \otimes \psi_-$ represents the tensor product of functions ψ_+ and ψ_- .

- (a) Express state Ψ as a linear combination of basis functions Ψ_{00} , Ψ_{01} , Ψ_{10} and Ψ_{11} , and determine the probability that each of these states will be registered if a measurement is performed on both particles.

(b) Suppose we now define an operator

$$\hat{W} = \sqrt{2} \left(\hat{I} \otimes \hat{A}_0 - \hat{A}_1 \otimes \hat{I} \right) \quad (14)$$

where \hat{I} is the identity operator, and \hat{A}_0 and \hat{A}_1 are the operators that were introduced in Problem 4. Find the function Φ that is obtained when \hat{W} is applied to Ψ , and determine whether this function can be expressed as

$$\Phi = \psi_a \otimes \psi_b \quad (15)$$

where ψ_a and ψ_b have the general form

$$\psi_a = \alpha_0 \psi_0 + \alpha_1 \psi_1 \quad (16)$$

and

$$\psi_b = \beta_0 \psi_0 + \beta_1 \psi_1 \quad (17)$$

If so, indicate what functions ψ_a and ψ_b look like explicitly.

- (c) Construct a quantum circuit that will produce state Φ using a combination of Hadamard and C-NOT gates.
- (d) Repeat parts (b) and (c) in case operator \hat{W} is defined as

$$\hat{W} = \sqrt{2} \left(\hat{A}_1 \otimes \hat{A}_0 - \hat{A}_0 \otimes \hat{A}_1 \right) \quad (18)$$